

1. ANALYSIS QUALIFYING EXAM, SPRING 2004

Instructions: Clearly explain and justify your answers. You may cite theorems from textbooks or that were proved in class as long as they are not what the problem explicitly asks you to prove. You may also use the results of prior problems or prior parts of the same problem when solving a problem – this is allowed even if you were unable to prove the previous results. Make sure to state the results that you are using and be sure to verify their hypotheses. All problems have **equal** value.

Notation: Let m denote Lebesgue measure on $(\mathbb{R}, \mathcal{B}_{\mathbb{R}})$ and (X, \mathcal{M}, μ) denote a finite measure space. Also for a bounded measurable function, $f : X \rightarrow \mathbb{C}$, let $M_f : L^2(\mu) \rightarrow L^2(\mu)$ denote the multiplication operator, $M_f h := fh$ for all $h \in L^2(\mu)$.

Exercise 1.1. In this problem $(X, \|\cdot\|)$ is an **infinite** dimensional normed space. Determine which of the following statements are true. For the true statements give a **brief** reason and for the false statements give a counter example.

- (a) If $A = \mathbb{R} \setminus E$ and $m(E) = 0$, then $\bar{A} = \mathbb{R}$.
 - (b) Every m – null set $E \in \mathcal{B}_{\mathbb{R}}$ is nowhere dense in \mathbb{R} .
 - (c) Every proper subspace $E \subset X$ is nowhere dense.
 - (d) If E is a subspace of X with non-empty interior, then $E = X$.
-

Exercise 1.2. Compute the values of the following two expressions. (You must justify your answers.).

(a)

$$\sum_{n=0}^{\infty} \int_0^{\infty} e^{-2x} \frac{(-1)^n}{(2n+1)!} x^{2n+1} dx$$

(b)

$$\int_0^{\infty} \left(\int_0^{\infty} x^2 e^{-x^2} \sin(x^2) e^{-yx} dx \right) dy.$$

You may find the following integration formula useful;

$$\int e^{-ax} \sin x dx = -\frac{1}{a^2 + 1} e^{-ax} [\cos x + a \sin x] + C.$$

Exercise 1.3. Suppose that $\{u_n\}_{n=1}^{\infty}$ is an orthonormal subset of Hilbert space, H , and S is a dense subset of H . Show $\{u_n\}_{n=1}^{\infty}$ is an orthonormal basis for H if

$$(1.1) \quad \|f\|_H^2 = \sum_{n=1}^{\infty} |\langle f | u_n \rangle|^2 \text{ for all } f \in S.$$